

1. 1D ionization. This problem is from the January, 2007, prelims. Consider a non-relativistic mass  $m$  particle with coordinate  $x$  in one dimension that is subject to an attractive delta-function potential at  $x = 0$ , i.e., a potential  $V(x) = -V_0\delta(x/a)$ , with  $V_0 > 0$ .

- (a) The ground state of the particle is a bound state. Find its wave function and binding energy.
- (b) The particle is now perturbed by a weak time-dependent potential  $V(x, t) = Fx \cos \omega t$ . Find the transition rate from the bound state to the continuum. (It may help to confine the particle in a large box  $|x| < L/2$  and take the limit  $L \rightarrow \infty$ .)

2. Spherical Square Potential. Consider low energy scattering of a particle of mass  $m$  from a spherical potential of radius  $a$ :

$$V(r) = \begin{cases} V_0 & r < a \\ 0 & r > a \end{cases},$$

where  $V_0$  may be either positive or negative.

- (a) Calculate the  $s$ -wave phase shift for incident energy  $E$ . Note that low energy scattering means  $ka \ll 1$ .
- (b) Can the  $s$ -wave phase shift be a multiple of  $\pi$ ? What happens in this case? Hint: Google "Ramsauer Effect."

3. A really, really square potential! This problem appeared on the May, 2004 prelims. A beam of particles of mass  $m$  and energy  $E$  propagates along the  $z$  axis of a coordinate system and scatters from the cubic potential

$$V = \begin{cases} v & \text{if } |x| \leq L, |y| \leq L, \text{ and } |z| \leq L, \\ 0 & \text{otherwise} \end{cases}$$

where  $v$  is a small constant energy.

- (a) Use the Born approximation to find an explicit formula for the scattering cross section  $\sigma = \sigma(\theta, \phi)$  as a function of the angles  $\theta$  and  $\phi$ . Recall that spherical coordinates of a point in space are related to the Cartesian coordinates  $(x, y, z)$  by  $x = r \sin \theta \cos \phi$ ,  $y = r \sin \theta \sin \phi$ , and  $x = r \cos \theta$ . The Born approximation is easy to evaluate in one coordinate system and hard in the other.
- (b) Under what circumstances is this approximation for the scattering cross section valid? Explain.

4. Neutron capture. (Based on a problem in Dicke and Witke, *Introduction to Quantum Mechanics*.) For a particular nucleus, the neutron absorption cross section, for 0.1 eV neutrons, is  $\sigma_a = 2.5 \times 10^{-18} \text{ cm}^2$ . What are the upper and lower bounds on the 0.1 eV neutron elastic scattering cross section?